Incremental Versus Conventional Hemodialysis for Preservation of Residual Kidney Function in Patients on Regular Hemodialysis

Hosam Ezzat Ebrahim^{1,*} MSc; Emad Allam Mohamed² MD ; Mohamed Ahmed El Sayed Ahmed² MD
Osama Hassan Bakheet² MD

*Corresponding Author:

Hosam Ezzat Ebrahim

cristiano9966@gmail.com

Received for publication October 31, 2020; Accepted January 3, 2021;
Published online January 3, 2021.

Copyright 2020 The Authors published by Al-Azhar University, Faculty of Medicine, Cairo, Egypt. All rights reserved. This an open-access article distributed under the legal terms, where it is permissible to download and share the work provided it is properly cited. The work cannot be changed in anyway or used commercially.

doi: 10.21608/aimj.2021.45156.1338

¹Clinical pathology Department, Military Medical Academy, Cairo, Egypt.

²Internal medicine and nephrology Department, Faculty of Medicine, Al-Azhar University, Cairo, Egypt.

Disclosure: The authors have no financial interest to declare in relation to the content of this article. The Article Processing Charge was paid for by the authors.

Authorship: All authors have a substantial contribution to the article.

ABSTRACT

Background: Incremental hemodialysis (HD) has several potential advantages for patients, physicians and health care systems. In incident HD patients, conservation of residual kidney function "RKF" is significant and is correlated with several advantages, including survival of patient, improved quality of life, improved overall nutritional status and less anemia. The aim of this work was to evaluate incremental hemodialysis (twice/week) compared with conventional dialysis (thrice/week) for the conservation of the residual kidneyfunction among patients initiating regular hemodialysis.

Aim of work: The primary outcome of the study was an evaluation of incremental hemodialysis (twice/week) compared with conventional dialysis (thrice/week) for the conservation of the residual kidney function among patients initiating regular hemodialysis.

Patient and Methods: We assigned 40 patients with chronic kidney disease stage V D who early started hemodialysis into 2 groups: Group 1: 20 patients underwent incremental hemodialysis (twice/week), and Group 2: 20 patients underwent conventional hemodialysis (thrice/week). All patients followed for laboratory findings of (complete blood picture, creatinine, urea, calcium, albumin, phosphorus, parathyroid hormone, alkaline phosphatase, glomerular filtration rate). Also, Kt/v at start and every month was done. Clinical outcome measures included; mortality, cardiovascular outcome, hospital admission and dialysis complications.

Results: In the studied population, the mean age was (45.95 ± 2.97) years. Regarding sex 60% were males and 40% were females. Regarding outcome data, (20%) of patients had hypotension, (15%) had hospital admission, while nobody suffered mortality. Comparative study among the two groups showed a substantial decrease in the incidence of hypotension in the incremental group (40 %) relative to the conventional group (10%) (p = 0.03).

Conclusion: To conclude, our research investigates the correlation among the frequency of HD treatment and survival of patient. In line with recent literature, our results show that some selected patients with acceptable RKF, sufficient control of interdialytic gaining weight, and low or moderate burden of comorbid disease (CCI < 5) in an incident HD population, an HD incremental approach may be deemed as an appropriate alternative to conventional thrice-weekly HD.

Keywords: Incremental dialysis; Conventional dialysis; Residual Kidney Function.

INTRODUCTION

There is substantial residual kidney function (RKF) for many HD-initiated patients. More than 90% of new ESRD hemodialysis patients start a standardized HD prescription 3 times a week in the United States¹.

Incremental HD twice-weekly regimens are deemed suboptimal, whereas in some, with restricted resources, this is the only choice. In peritoneal dialysis, maintaining residual urine production is a significant goal, but this issue is not typical in hemodialysis. Vilar et al. and Fernández-Lucas et al. both found that patients beginning with twice-weekly incremental HD had better survival and greater residual urine production than patients beginning with a thrice-weekly regimen².

Incremental dialysis uses the principle of changing the dialysis dose as per to RRF so that the dialysis dose is individualized. The basis is to provide adequate dialysis to achieve supra-minimum elimination of uremic solutes and the control of hypervolemia and then to increase the dialysis dose as the RRF decreases. In order to provide complete composite clearance, the general idea is to measure the total amount of urea extraction during dialysis and add it to the residual renal urea clearance (KRU)^{3,4}.

Incremental HD offers patients, physicians, and health systems several possible advantages. In incident HD patients, 'preservation of RKF' is

significant and is correlated with many advantages, like patient longevity, improved quality of life, enhanced overall dietary status and decreased anemia (The Importance of Residual Kidney Function, see above). The "longevity of vascular access" associated with less regular arteriovenous fistula or graft cannulations are another advantage of incremental HD. In an examination from the FHN study, more frequent HD decreased the composite endpoint of vascular access loss, repair or access-related hospitalization. The chance of a first access event was 76 % higher with regular HD than with conventional HD⁵.

For less frequent HD therapy regimens, economic advantages also need to be considered. Conventional thrice-weekly HD treatments cost around \$89,000 per person per year in the United States, with a gross annual cost of \$42 billion (\$34 billion covered by Medicare, the rest by , private insurance, Medicaid or out-of-pocket payments)⁶.

PATIENTS AND MATERIALS

We assigned 40 patients with chronic kidney disease stage V D who early started hemodialysis into 2 groups: Group 1: 20 patients underwent incremental hemodialysis (twice/week). and Group 2: 20 patients underwent conventional hemodialysis (thrice/week).

During 6 months in Nephrology department, Military Hospitals (AL-maadi, kobry AL-koba and air force military hospital).

All patients will be subjected to: History & complete physical examination, Routine laboratory investigations: CBC, liver and kidney function Calcium, phosphorus, intact parathyroid hormone GFR and KT/V. Clinical outcome

Measures: Mortality, Cardiovascular outcome, Hospital admission and Dialysis complications,

Statistical analysis:

Using MedCalc ver. 18.2.1 (MedCalc, Ostend, Belgium), data input, processing and statistical analysis have been conducted out. Significance tests were used (Mann-Whitney's, Chi square, tests, factorial ANOVA, logistic regression analysis, and analysis of the ROC Curve). Data were supplied and appropriate analysis was carried out as per type of data obtained for each variable (parametric and non-parametric). P-values of lower than 0.05 (5%) were deemed statistically important.

RESULTS

In the studied population, the mean age of all patients was (45.95 ± 2.97) years, and the average UOP was (1124.3 ± 337.2) ml/day. Regarding gender of the patients, the majority (60%) of patients were males; while (40%) were females, with (65%) had DM, and (75%) had HTN.

Regarding outcome data, (20%) of patients had hypotension, (15%) had hospital admission, while nobody suffered mortality.

Comparative studies: The 40 CKD patients were split into two separate groups accordingly to the dialysis

technique: Conventional group (20 patients) and Incremental group (20 patients).

Non-significant differences in all basic clinical data were observed in a comparative analysis among the two groups ($p > 0.05$).

Non-significant differences in all baseline laboratory data were observed in the comparative analysis among the two groups ($p > 0.05$).

ROC curve analysis to predict patients (6-months) improvement (Table 1, Fig. 1 - 4): By using ROC-curve analysis, Incremental dialysis technique expected patients with hemoglobin improvement, with excellent (93%) accuracy, sensitivity= 75% and specificity= 100% ($p < 0.05$).

By using ROC-curve analysis, Incremental dialysis technique predicted patients with platelets improvement, with fair (70%) accuracy, sensitivity= 55% and specificity= 100% ($p < 0.05$).

By using ROC-curve analysis, Incremental dialysis technique predicted patients with GFR improvement, with perfect (100%) accuracy, sensitivity= 100% and specificity= 100% ($p < 0.05$).

By using ROC-curve analysis, Incremental dialysis technique predicted patients with uric acid improvement, with fair (70%) accuracy, sensitivity= 50% and specificity= 90% ($p < 0.05$).

By using ROC-curve analysis, Incremental dialysis technique expected patients with Ca improvement, with excellent (96%) accuracy, sensitivity= 100% and specificity= 80% ($p < 0.05$).

By using ROC-curve analysis, Incremental dialysis technique expected patients with Ph improvement, with excellent (94%) accuracy, sensitivity= 85% and specificity= 90% ($p < 0.05$).

By using ROC-curve analysis, Incremental dialysis technique expected patients with PTH improvement, with excellent (91%) accuracy, sensitivity= 80% and specificity= 90% ($p < 0.05$).

By using ROC-curve analysis, Incremental dialysis technique predicted patients with albumin improvement, with fair (79%) accuracy, sensitivity= 55% and specificity= 90% ($p < 0.05$).

By using ROC-curve analysis, Incremental dialysis technique predicted patients with Kt/V improvement, with good (89%) accuracy, sensitivity= 60% and specificity= 100% ($p < 0.05$).

By using ROC-curve analysis, Incremental dialysis technique showed non-significant predictive values regarding TLC, creatinine, urea, Na and K improvements ($p > 0.05$).

Variable	AUC	SE	Sensitivity (%)	Specificity (%)	P value
Hb	0.930	0.038	75	100	<0.0001**
PLT	0.700	0.087	55	100	0.022*
TLC	0.585	0.093	65	60	0.3652
Creat. (post-dialysis)	0.660	0.090	90	60	0.0781
Urea (post-dialysis)	0.600	0.094	100	40	0.2882
GFR	1.000	0	100	100	<0.0001**
Uric acid	0.703	0.083	50	90	0.015*
Na	0.623	0.096	40	100	0.2053
K	0.588	0.091	90	30	0.3399
Ca	0.968	0.023	100	80	<0.0001**
Ph	0.941	0.033	85	90	<0.0001**
PTH	0.915	0.043	80	90	<0.0001**
Alb.	0.790	0.072	55	90	0.0001**
Kt/V	0.895	0.048	60	100	<0.0001**

ROC (Receiver operating characteristic), AUC= Area under curve, SE= Standard Error.

Table 1: Roc-curve of Incremental dialysis technique to predict patients (6-months) improvement:

Fig.1: ROC curve of hemoglobin and platelets (Incremental dialysis technique).

Fig.2: ROC curve of GFR and uric acid (Incremental dialysis technique).

Fig.3: ROC curve of Ca and Ph (Incremental dialysis technique).

Fig.4: ROC curve of PTH, albumin and Kt/V (Incremental dialysis technique).

Variable		Conventional group (20)	Incremental group (20)	Chi square test
				P value
Hypotension	+ve	8 (40%)	2 (10%)	= 0.03*
Hospital admission	+ve	6 (30%)	2 (10%)	= 0.1185

Table 2: Comparison between the two groups with respect to the results data using the Chi square test:

Comparative among the two groups identified; substantial decrease in the incidence of hypotension in incremental groups (40 %); relative to conventional groups (10%); with a very significant statistical difference ($p = 0.03$) (Table 2, Fig. 5).

Non-significant differences in hospital admission were observed in the comparative analysis among the 2 groups ($p > 0.05$).

Fig.5: Comparison between the 2 groups of patients regarding hypotension.

DISCUSSION

This was a prospective comparative research performed in 40 patients with chronic kidney disease (CKD); to assess incremental hemodialysis (twice/week) compared with conventional dialysis (thrice/week) for the conservation of residual kidney function among patients initiating regular hemodialysis.

The study will be conducted in patients with chronic end stage kidney disease through 2018 and 2019 in Military hospitals (Maadi Military Hospital, kobriekoba military hospital and air force military hospital). We assigned 40 patients with chronic kidney disease stage V D who early started hemodialysis.

Group1: 20 patients underwent incremental hemodialysis (twice/week) and Group 2: 20 patients underwent conventional hemodialysis (thrice/week).

Patients have been followed for laboratory findings of (complete blood picture, serum creatinine, urea, calcium, albumin, phosphorus, parathyroid hormone, alkaline phosphatase, glomerular filtration rate).

Regarding Baseline data, we found that; the mean age of all patients was (45.95 ± 2.97) years, and the average UOP was (1124.3 ± 337.2) ml/day. Regarding gender of the patients, the majority (60%) of patients were males; while (40%) were females, with (65%) had DM, and (75%) had HTN, which came in agreement with A. Mathew et al.⁷ and Lucas et al.⁸.

Regarding Outcome data, (20%) of patients had hypotension, (15%) had hospital admission, while nobody suffered mortality, which came in agreement with Lucas et al.⁸

Lucas et al reported that in the 2- and 3-HD / week groups, hospitalization period was 4 (0-15) and 11 (2-14) days/patient-years, respectively (median and IQR, $P = 0, 068$).

No substantial difference in the comparative study between the 2 groups with respect to all basic clinical data ($p > 0.05$), which came in agreement with A. Mathew et al.⁷, Lucas et al.⁸ and Liu et al.⁹

A. Mathew et al. reported that, no statistical disparity with all-cause mortality in subgroups of age, gender

, ethnicity, central venous catheter use or diabetes condition in preset subgroup analyses of the matched cohort compared incremental HD to conventional HD.⁷

We found marked increase in UOP in incremental group; compared to conventional group; during the serial measurements ($p < 0.05$), which came in agreement with Liu et al.⁹, A. Mathew et al.⁷, Lucas et al.⁸, A. T. Mathew et al.¹⁰ and Merino et al.¹¹

Liu et al reported that those on incremental HD appeared to be female and thinner relative to those on traditional HD. The amount of urine in the incremental HD group was higher relative to the conventional HD group.⁹

Lucas et al reported that the daily volume of urine and GFR (glomerular filtration rate) is higher throughout the 2 HD / week group. In patients who began 2- and 3 HD / week, the GFR range was 2–12, 6 mL / min and 0, 2–13, 6 mL / min, respectively.⁸

We found marked increase in hemoglobin in incremental group; compared to conventional group; during the serial measurements ($p < 0.05$), which came in agreement with Merino et al.¹¹ and A. Mathew et al.⁷

Hemoglobin and hematocrit enhanced after 1 year and the mean dose of erythropoiesis stimulating factors (ESF) did not raise significantly, reported by Merino et al. 2017, which was darbepoetin (mcg / week), 27 ± 21 and 30 ± 22 mcg / week, $p = 0.725$ in our case.¹¹

Liu et al reported that those on incremental HD appeared to be female and thinner relative to patients on traditional HD. The calcium and hemoglobin levels between the groups were similar.⁹

Lucas et al reported that both groups had similar hemoglobin concentrations; however, the weekly erythropoietin dose was substantially lower in patients, who began 2 HD / week.⁸

We found marked increase in post-dialysis urea in incremental group; compared to conventional group; during the serial measurements ($p < 0.05$), which came in agreement with A. Mathew et al. 2016.⁷

A. Mathew et al reported that patients with an incremental HD regimen and those with sufficient baseline RKF had substantially more conservation of renal clearance of urea in that study.⁷

We found marked increase in GFR in incremental group; compared to conventional group; during the serial measurements ($p < 0.05$), which came in agreement with Kalantar-Zadeh et al.¹², Bolasco et al.¹³, Rhee et al.¹⁴, Liu et al.⁹, A. Mathew et al.⁷, Lucas et al.⁸ and Obi et al.¹⁴

In reality, Bolasco et al recorded faster loss of RKF by twice-weekly or more regular hemodialysis as a negative predictive factor for both mortality and morbidity; however, nephrologists have long undervalued the clinical significance of RKF.¹³

A. Mathew et al reported that, a prospective study of 168 patients with HD incidence showed that in twice-weekly HD-initiated patients, the proportion of

patients with RKF loss was substantially lower in twice-weekly HD-initiated patients relative to conventional thrice-week HD-initiated patients⁷.

Obi et al reported that both the clearance of renal urea and the amount of urine in incremental vs conventional regimens ($P = 0.001$ for both) displayed considerably slower decreases over time in this matched cohort.¹⁴

We found marked increase in calcium in incremental group; compared to conventional group; during the serial measurements ($p < 0.05$), which came in agreement with A. Mathew et al.⁷ and Obi et al.¹⁴

A. Mathew et al reported that, the final cohort of the study consisted of 87,718 patients from 1737 facilities, comprising 201 frequent (\$4 times weekly) HD patients from 158 facilities and 682 incremental (twice weekly or less) HD patients from 444 facilities, with increased calcium level in incremental group 9.1 ± 0.5 .⁷

Obi et al reported that, Trends in the selected parameters indicated lower dialysis frequency, shorter dialysis period, lower hemoglobin and corrected serum calcium concentrations in patients treated with an incremental regimen.¹⁴

Liu et al reported that, there are no statistically substantial variations in serum levels of calcium, phosphate, albumin, hemoglobin or hospitalization rates among groups. Serum calcium (SMD = -0.397, 95% CI: -0.523 to -0.272, $P < 0.001$; I² = 0 %) relative to laboratory values at the end of follow-up (subgroup 1).⁹

We found marked decrease in phosphorus in incremental group; compared to conventional group; during the serial measurements ($p < 0.05$), which came in agreement with A. Mathew et al.⁷, Merino et al.¹¹ and Liu et al.⁹

A. Mathew et al reported that, the final cohort of the study consisted of 87,718 patients from 1737 facilities, comprising 201 frequent (\$4 times weekly) HD patients from 158 facilities and 682 incremental (twice weekly or less) HD patients from 444 facilities, with decreased phosphorus in incremental group 4.3 ± 1.0 .⁷

Merino et al reported that after 1 year, serum phosphate level control had improved, but the variations were not statistically relevant.¹¹

Liu et al reported that, Serum phosphate levels in the incremental HD group were lower, and serum albumin levels were higher.⁹

We found marked decrease in PTH in incremental group; compared to conventional group; during the serial measurements ($p < 0.05$), which came in agreement with Merino et al.¹¹

Merino et al reported that after 1 year, control of PTH levels improved, but the variations were not statistically relevant.¹¹

We found marked decrease in albumin in incremental group; compared to conventional group; during the serial measurements ($p < 0.05$), which came in disagreement with Liu et al.⁹

Liu et al reported that, Serum phosphate levels in the incremental HD group were lower, and serum albumin levels were higher.⁹

We found marked decrease in Kt/V in incremental group; compared to conventional group; during the serial measurements ($p < 0.05$), which came in agreement with Lucas et al.⁸

A comparative study among the 2 groups demonstrated; a substantial decrease in the incidence of hypotension in the incremental group (40%); a highly substantial statistical difference ($p = 0.03$), relative to the conventional group (10%); which came in agreement with A. Mathew et al.⁷, Merino et al.¹¹, Kalantar-Zadeh et al.¹² and A. T. Mathew et al.¹⁰

Merino et al reported that, the average baseline systolic blood pressure (mmHg) was 145 ± 12 mmHg versus 151 ± 14 after 12 months ($p = 0.165$) and the average baseline diastolic blood pressure was 72 ± 13 versus 67 ± 20 at baseline and 1 year ($p = 0.243$). Blood pressure was similar at baseline relative to 12 months.¹¹

Comparative study among the two groups showed no substantial difference in hospital admission ($p > 0.05$), which came in agreement with A.T. Mathew et al.¹⁰, Liu et al.⁹ and Lucas et al.⁸

A. T. Mathew et al reported that, more frequent HD decreased the composite endpoint of vascular access loss, repair, or access-related hospitalization in the analysis of the FHN study.¹⁰

Kalantar-Zadeh et al reported that, this incremental approach to hemodialysis offers a more gradual and tolerable transition to renal replacement treatment, possibly resulting in improved quality of life linked to health, longer conservation of residual kidney function, reduction of inflammatory and oxidative stress caused by hemodialysis, reduced rate of intra-dialytic hypotension, decreased stimulating agent dose of erythropoiesis and decreased mortality and morbidity.¹²

CONCLUSION

To conclude, our research investigates the correlation among the frequency of HD treatment and survival of patient. In line with recent literature, our results show that some selected patients with acceptable RKF, sufficient control of interdialytic gaining weight, and low or moderate burden of comorbid disease (CCI < 5) in an incident HD population, an HD incremental approach may be deemed as an appropriate alternative to conventional thrice-weekly HD.

REFERENCES

1. O'Hare, A.M., Wong, S.P., Margaret, K.Y., Wynar, B., Perkins, M., Liu, C.-F., et al. Trends in the timing and clinical context of maintenance dialysis initiation. *Journal of the American Society of Nephrology*. 2015; 26 (8), 1975–81.
2. Vilar, E., Wellsted, D., Chandna, S.M., Greenwood, R.N., and Farrington, K. Residual renal function improves outcome in incremental haemodialysis

- despite reduced dialysis dose. *Nephrology Dialysis Transplantation*. 2009; 24 (8), 2502–10.
3. Wong, J., Vilar, E., Davenport, A., and Farrington, K. Incremental haemodialysis. *Nephrology Dialysis Transplantation*. 2015; 30 (10), 1639–48.
 4. Kessler, M., Canaud, B., Pedrini, L.A., Tattersall, J., ter Wee, P.M., Vanholder, R., et al. European Best Practice Guidelines for Haemodialysis (part 1): Section II: Haemodialysis adequacy. *Nephrol Dial Transpl*. 2002; 17 16–31.
 5. Suri, R.S., Larive, B., Sherer, S., Eggers, P., Gassman, J., James, S.H., et al. Risk of vascular access complications with frequent hemodialysis. *Journal of the American Society of Nephrology*. 2013; 24 (3), 498–505.
 6. Collins, A.J., Foley, R.N., Gilbertson, D.T., and Chen, S.-C. United States Renal Data System public health surveillance of chronic kidney disease and end-stage renal disease. *Kidney International Supplements*. 2015; 5 (1), 2–7.
 7. Mathew, A., Obi, Y., Rhee, C.M., Chen, J.L., Shah, G., Lau, W.-L., et al. Treatment frequency and mortality among incident hemodialysis patients in the United States comparing incremental with standard and more frequent dialysis. *Kidney International*. 2016; 90 (5), 1071–9.
 8. Lucas, M.F., Teruel, J.L., Ruíz-Roso, G., Díaz, M., Raoch, V., Caravaca, F., et al. Incremental hemodialysis schedule in patients with higher residual renal function at the start of dialysis. *Advances in Nephrology*. 2014.
 9. Liu, Y., Zou, W., Wu, J., Liu, L., and He, Q. Comparison between incremental and thrice-weekly haemodialysis: Systematic review and meta-analysis. *Nephrology*. 2019; 24 (4), 438–44.
 10. Mathew, A.T., Obi, Y., Rhee, C.M., Chou, J.A., and Kalantar-Zadeh, K. Incremental dialysis for preserving residual kidney function—Does one size fit all when initiating dialysis? in: *Seminars in Dialysis, Wiley Online Library*, pp. 2018; 343–52.
 11. Merino, J.L., Domínguez, P., Bueno, B., Amézquita, Y., Espejo, B., and Paraíso, V. Application of model of incremental haemodialysis, based on residual renal function, at the initiation of renal replacement therapy. *Nefrología (English Edition)*. 2017; 37 (1), 39–46.
 12. Kalantar-Zadeh, K., Crowley, S.T., Beddhu, S., Chen, J.L., Daugirdas, J.T., Goldfarb, D.S., et al. Renal replacement therapy and incremental hemodialysis for veterans with advanced chronic kidney disease. in: *Seminars in Dialysis, Wiley Online Library*, pp. 2017; 251–61.
 13. Bolasco, P., Cupisti, A., Locatelli, F., Caria, S., and Kalantar-Zadeh, K. Dietary management of incremental transition to dialysis therapy: once-weekly hemodialysis combined with low-protein diet. *Journal of Renal Nutrition*. 2016; 26 (6), 352–9.
 14. Obi, Y., Streja, E., Rhee, C.M., Ravel, V., Amin, A.N., Cupisti, A., et al. Incremental hemodialysis, residual kidney function, and mortality risk in incident dialysis patients: a cohort study. *American Journal of Kidney Diseases*. 2016; 68 (2), 256–5.