

Impact Of Royal Jelly and L-Carnitine on Improving the Histological Changes of Cyclophosphamide-Induced Testicular Cytotoxicity in Adult Male Albino Rats

Khalifa Abd-Elrazik Khalifa Swidan,¹ MSc, Gamal Soliman Hassan Al-Gharabawi,² MD, Tamer M.M. Abu Emara,² MD.

* Corresponding Author:

Khalifa Abd-Elrazik Khalifa Swidan
dr.khalifa.com@gmail.com

Received for publication
February 05, 2021; Accepted
July 17, 2022; Published online
July 17, 2022.

doi: 10.21608/aimj.2022.115985.1791

Citation: Khalifa A. , Gamal S. , Tamer M.. Impact Of Royal Jelly and L-Carnitine on Improving the Histological Changes of Cyclophosphamide-Induced Testicular Cytotoxicity in Adult Male Albino Rats. AIMJ. 2022; Vol.3-Issue7 .7-13.

¹Histology Department, Faculty of Medicine, Al-Azhar University, Assiut, Egypt

²Histology Department, Faculty of Medicine, Al-Azhar University, Cairo, Egypt.

ABSTRACT

Background: Cyclophosphamide is anticancer chemotherapeutic drug and immunosuppressive agent for organ transplantation. Wide range of adverse effects including reproductive toxicity demonstrated following its administration.

Aim of the study: To evaluation the impact of royal jelly and L-Carnitine on improving the histological and Immunohistochemical changes of experimentally Cyclophosphamide-induced testicular cytotoxicity in adult male albino rats.

Materials and Methods: forty adult male albino rats Were divided into two groups, the control group1 (ten rats) and treated group 2 (thirty rats) subdivided into three equal subgroups (a, b and c). Subgroup (a): Ten rats treated with cyclophosphamide. Subgroup (b): Ten rats given royal jelly with cyclophosphamide. Subgroup (c): Ten rats given L-Carnitine and cyclophosphamide. The testes excised and prepared for light microscopic examination and stained with H&E and Masson's trichrome stains and immunohistochemically for detecting bcl-2 proteins.

Results: Seminiferous tubules distorted and shrunken with marked reduction in the thickness of the germinal epithelium that appeared detached from irregular thick basement membrane and wide empty lumen devoid of spermatozoa, increased amount of interstitial tissue, contain congested blood vessels and covered by thick tunica albuginea. Royal jelly and L-Carnitine had a potent protective effect against the testicular toxicity and showed a notable effect.

Conclusion: Cyclophosphamide produced marked changes in the histological structure of the. Royal jelly and L-Carnitine have a potent protective effect against the toxicity.

Keywords: Cyclophosphamide; royal jelly; L-Carnitine; bcl-2.

Disclosure: The authors have no financial interest to declare in relation to the content of this article. The Article Processing Charge was paid for by the authors.

Authorship: All authors have a substantial contribution to the article.

Copyright The Authors published by Al-Azhar University, Faculty of Medicine, Cairo, Egypt. Users have the right to read, download, copy, distribute, print, search, or link to the full texts of articles under the following conditions: Creative Commons Attribution-Share Alike 4.0 International Public License (CC BY-SA 4.0).

INTRODUCTION

Survival rates from most childhood cancers have increased dramatically in recent decades, with chemotherapy a mainstay of treatment for the majority of these patients. Ideally, chemotherapeutic agents would specifically target cancerous cells with many off-target effects of cytotoxic treatments, including on the gonads.¹ Cyclophosphamide (CP) is a drug belongs to the class of oxazaphosphorines, which is essential anticancer chemotherapeutic drug, immunosuppressive for organ transplantation and

many acute and chronic benign diseases as systemic lupus erythematosus, multiple sclerosis, and nephritic syndrome. Also effectively maintained remission of

childhood nephrotic syndrome with frequent relapses.² A range of adverse effects of CP have been reported including reproductive toxicity, as diminished sperm counts and absence of spermatogenic cycles in their testicular tissue.³ Many genes, such as pro-apoptotic member (Bax) and anti-apoptotic member (Bcl-2 and Bcl-w), regulate the apoptosis process.⁴

Royal jelly is one of honey products. It stimulates cell survival, cell growth and cell differentiation; it has anti-tumor and anti-metastatic effect.⁵ L-carnitine (LC) found in epididymis and sperm, its concentration is about 2000 times higher than in plasma, it is important in lipid metabolism that

facilitates long-chain fatty acids' β -oxidation in mitochondria and is necessary for energy production.⁶ It acts as a substantial non-enzymatic antioxidant that protects the cell mitochondrial membrane, and DNA integrity against free oxygen radicals. Several studies performed to evaluate the effect of carnitine on infertile men, indicating the improvement of sperm fertility, count, and motility, so it is effective in improving fertility⁷.

This work aimed to investigate the impact of royal jelly and L-Carnitine on improving the histological, Immunohistochemical changes of experimentally Cyclophosphamide-induced testicular cytotoxicity.

MATERIAL AND METHODS

Forty rats used, divided into two groups; control group (1): ten rats injected intraperitoneal with normal saline. Second group: subdivided into three subgroups (a, b and c). (a): ten rats injected intraperitoneally with cyclophosphamide solution at a dose of 50 mg/kg/day in every alternative day for 14 days. (b): ten rats given royal jelly orally at dose of 1000 μ g/kg/day for 1 week before and with the use of cyclophosphamide at a same dose with second group. (c): ten rats given L carnitine intra-peritoneally at daily dose of 500 mg/kg/day for 1 week before and with the use of cyclophosphamide at a same dose with second group. Testes excised and put in Bouin's fixative. Sections (5um thick) obtained and stained with H&E, Masson's trichrome stains and immunohistochemically for localization of bcl-2 proteins.

Image analysis and morphometric study:

Stained sections were morphometrically analyzed using image analyzer computer system to detect the significance of changes in area% of collagen fibers, area% and optical density for bcl-2.

Statistical analysis:

The data analyzed by using SPSS version 23, expressed as means and standard deviation (SD) then analyzed statistically using one-way analysis of variance (ANOVA) for comparison between the different groups with p value less than 0.05 (the level of significance).

RESULTS

Group I: Testicular tissue formed of regular rounded or oval shaped sections of seminiferous tubules, surrounded by a regular thin basement membrane, and lined by seminiferous epithelium formed of spermatogenic cells at different stages of maturation with lumen full of spermatozoa (Fig. 1a). The interstitial tissue contains thin-walled blood vessels, cells of Leydig covered by a fibrous tunica albuginea (Fig. 2a). Group IIa: Seminiferous tubules markedly distorted with irregular outlines, smaller in diameters than control and show many vacuoles and the lumen is devoid of spermatozoa (Fig.1b). The amount of

interstitial areas increased and contain some thick walled congested blood vessels and increased number of cells of Leydig that arranged in small clumps and covered by a thick tunica albuginea (Fig. 2b). Group IIb: The majority of the seminiferous tubules appeared regular in their outlines and architecture with the appearance of the germinal cell populations including different types of germ cells with some vacuoles and the lumen contain spermatozoa (Fig. 1c). Interstitial tissue, similar to the control group, contains thin-walled blood vessels, cells of Leydig, covered by a thin fibrous tunica albuginea (Fig. 2c). Group IIc: Regular seminiferous tubules, normal architecture with the appearance of the germinal cell populations with lumen full of spermatozoa (Fig. 1d). Interstitial tissue similar to that of control group, thick walled blood vessels in some sections, covered by a thick tunica. (Fig. 3d).

Immunohistochemical study: (1) group I: Strong immune reactivity of bcl-2 protein in the spermatids, primary spermatocytes, however moderate reactivity in spermatogonia (fig.3a). Group IIa: Negative immune reactivity in spermatogonia with minimal immune reactivity in spermatids and primary spermatocytes (fig.3b). Group IIb: Strong expression of bcl-2 protein in the spermatids and primary spermatocytes, Spermatogonia showed moderate expression. (fig.3c). Group IIc: it was similar to (group IIb) (fig.3d).

Morphometrical and statistical results:

1. Area % of collagen fibers: Significant increase ($P < 0.001$) in CP only treated (group IIa) in relation to control (group 1). Also, significant increase ($p < 0.01$) in (group IIb) and (group IIc) in relation to (group 1) and a significant decrease in relation to (group IIa). However, there was a non-statistically significant difference ($p > 0.05$) in (group IIb) in relation to (group IIc). (Table.1 & Fig. 4).

2. Area % of bcl-2 expression in the germinal epithelium: There was a significant decrease of expression in the germinal epithelium of seminiferous tubules ($P < 0.001$) in (group IIa) in relation to (group 1). Also, there was a significant decrease in bcl-2 expression ($p < 0.01$) in (group IIb) and (group IIc) in relation to (group 1), and a significant increase in relation to (group IIa). However, there was a non-statistically significant difference ($p > 0.05$) in (group IIb) in relation to (group IIc). (Table2, Fig. 5).

DISCUSSION

Testis is considered the most important organ in male reproductive system. It is characterized by two main functions, synthesis of steroid hormones and production of spermatozoa. Various chemical factors affect spermatogenesis such as drugs and toxic elements.⁸ Spermatogenesis is the sophisticated process of sperm formation and maturation. It takes place in the seminiferous epithelium and is represented by continuous germ cell maturation towards the center, high incidence of apoptosis occurs to discard the excessive germ cells.⁹ Obeyes and Mahood, reported that the testicular dysfunction

is a common long term sequel of cytotoxic drugs used in the treatment of many malignancies⁸.




Fig. 1: Group (a): Seminiferous tubules (ST) lined by well-arranged epithelium have lumen (L) containing sperms with interstitial tissue (IT) showing Leydig cells (Lc) and normal blood vessel (Bv). Group (b): (ST) containing vacuoles (V), (L) devoid of sperms. (IT) showing (Lc), thick walled (Bv). Group (c): (ST) lined by well-differentiated epithelium and have (L) containing sperms and some (V). (IT) showing (Lc). Group (d): (ST) lined by well-differentiated germinal epithelium, (L) full of spermatozoa. (Hx&E, X400)




Fig. 2: Group (a) (ST) surrounded by thin basement membrane (yellow arrow). (IT) containing normal (Bv), covered by a fibrous tunica albuginea (T). Group (b): (ST) surrounded by thick basement membrane, (IT) showing great amount of collagen fibers (red arrow) covered by thick (T). Group (c): (ST) with regular basement membrane (arrow), covered by a thin fibrous (T). Group (d): (ST) surrounded by thin, regular basement membrane (arrow), (IT) showing fair amount of collagen fibers (arrowhead) and covered by thick (T). (Masson's Trichrome Stain, X400).




Fig. 3: Group (a) Maximum expression of bcl2 in spermatids (red arrow) and primary spermatocytes (yellow arrows) with moderate immunoreactions in spermatogonia (arrow heads). group (b) Median expression in primary spermatocytes (arrows), minimal immunoreactions in spermatogonia (arrowhead). group (c) positive in spermatids (red arrow), primary spermatocytes (yellow arrow) and moderate reactivity in spermatogonia (arrowhead). group (d) positive reactivity in spermatids (red arrow), primary spermatocytes (yellow arrow) and moderate in spermatogonia (arrowhead). (Immunoperoxidase technique for bcl-2 X400).

Collagen fibers	Mean & SD	Median	Range	IQR	P
Control group	9.77 ± 2.176	10.30	6.10, 13.30	7.88, 11.13	< 0.001
Cyclophosphamide group	22.52 ± 3.024	22.10	18.70, 28.00	19.80, 24.92	
L-Carnitine group	14.16 ± 2.775	14.80	8.20, 17.70	13.27, 15.70	
Royal jelly group	16.52 ± 1.902	16.60	13.50, 20.30	15.23, 17.48	

Data is expressed as mean and standard deviation, median, range and interquartile range. P is significant when < 0.05.

Table 1: Comparison between the area percent of collagen fibers among the studied groups.




Fig. 4: Variations of the area percent of collagen fibers means among the studied groups.

Bcl-2 expression	Mean & SD	Median	Range	IQR	P
Control group	5.98 ± 1.570	6.13	3.85, 8.22	4.50, 7.29	< 0.001
Cyclophosphamide group	2.35 ± 1.393	2.49	0.57, 4.05	0.75, 3.71	
L-Carnitine group	4.05 ± 1.476	4.19	1.99, 5.88	2.46, 5.42	
Royal jelly group	4.09 ± 1.146	4.54	2.36, 5.65	2.81, 4.94	

Data is expressed as mean and standard deviation, median, range and interquartile range. P is significant when < 0.05.

Table 2: Comparison between the area percent of bcl-2 expression among the studied groups.




Fig. 5: Variations of the area percent of bcl-2 expression means among the studied groups.

Since the period of spermatogenic cycle in rats is 12 days; the treated animals left 14 days in subgroup (a) and 21 days in subgroups (b) and (c), a period, which is more than one spermatogenic cycle, to achieve the morphological affection of cyclophosphamide treatment and protection by other agents.

Cyclophosphamide produced marked degenerative changes in the form of reduction of the diameter of the seminiferous tubules, distortion of their shapes, irregularity in outlines, thickening of their basement membrane and an arrest of the normal stages of development of their germinal epithelium with absence of mature sperms. Many vacuoles, more numerous towards the basement membrane, these findings were in agreement with those of Kamarzaman et al. Tunica albuginea was very thick as compared to control group confirmed by Masson's trichrome stain. These findings correlated with observations of Kanth et al.,¹¹. The tunica vasculosa showed severe congestion of blood vessels, findings similar to that of Oluwole et al.,¹² which might be due to hyperemia secondary to increased cell activity and metabolism.¹¹ The oxidative stress caused by cyclophosphamide results in generation of reactive oxygen species (ROS) which play a critical role in the initiation and progression of fibrotic diseases. Nicotinamide adenine dinucleotide phosphate (NADPH) oxidase is the predominant enzyme source for ROS generation and now recognized as a key

mediator of cell proliferation and matrix accumulation¹⁴.

When adding royal jelly to the treatment with cyclophosphamide for two weeks showed marked degree of recovery from the testicular damage and spermatogenic cells reduced to few discrete layers, so lumen of the tubules appeared wide as compared to control. Testis of this group showed the majority of the seminiferous tubules appeared regular in their outline, germinal epithelium consisted of spermatogenic cells and Sertoli cells. The newly formed spermatogonia were next to the basement membrane, while the advanced spermatogenic stages were very close to the lumina in which the spermatozoa appeared and tubules surrounded by a thin basement membrane covered externally by a thin tunica albuginea. The interstitial tissue stroma, containing normal blood vessels, these results were in agreement with Karaca et al.,¹⁵. The protective effect of royal jelly may be due to its components, such as antioxidants, vitamins A, E, C, D and B complex, these vitamins themselves had anticancer effect and protective effects against the genotoxicity of chemotherapy and radiotherapy¹⁶.

The treatment with cyclophosphamide and LC showed also marked degree of recovery; the majority of the seminiferous tubules appeared regular in their outline and the germinal epithelium consisted of

spermatogenic cells and spermatozoa appeared in the lumen. LC is a nutrient with antioxidant effects, which has pivotal roles in energy production as it facilitates the passage of fatty acids into the mitochondria matrix for oxidation¹⁷. Since the main source of energy in testicular cells is fatty acid oxidation, LC role in transporting fatty acids into the sperm mitochondrial matrix for energy production is essential.¹⁸

The expression of Bcl-2 was downregulated in testicular tissues from cyclophosphamide treated rats compared to healthy controls. Similar findings reported by Hoda et al., that the germ cell apoptosis predominantly occurred in spermatogonia and to less extent in spermatocytes.¹⁹ Group (IIa) revealed negative reactivity in spermatogonia, minimal reactivity in spermatids and primary spermatocytes, CP induced apoptosis in the spermatogenic cells, as indicated by the highly significant decrease in the expression of bcl-2 protein. CP initiate a series of events that end in apoptosis. The ROS are key elements in this apoptosis, the amounts and activities of ROS-scavenging enzymes, including peroxidase, ascorbate, catalase and superoxide dismutase are strongly reduced with CP. These data imply that CP-exposed cells lose their ability to scavenge ROS and this loss results in oxidative damage and cell apoptosis²⁰.

Group (IIb): Strong expression in the spermatids and primary spermatocytes, these findings were in agreement with Karaca et al, who studied the effects of (RJ) on testicular damage in streptozotocin (STZ)-induced diabetic rats¹⁵. Cyclophosphamide with LC were nearly similar to group IIb this was in agreement with Roy et al., who indicated that mice fed with LC showed increased PGC1 α levels in the testis, which probably regulates steroidogenesis by increasing Bcl2 mRNA.²¹ In addition, LC effectively prevents apoptosis by carrying the accumulated long chain fatty acid surrounding mitochondria in testes of diabetic rats.²² These findings were in agreement with Vardiyan R et al, who found that an increase in bcl-2 expression and a decrease in Bax expression in mice administered both formalin and L-carnitine²³. Mohammadi V et al, found out that LC successfully increased expression of bcl-2 gene in rooster's testis²⁴. Finally, oxidative stress leads to an increase in the release of cytochrome C from the mitochondria, which activates the apoptotic mechanism in the cells by activating caspases (caspase-3/7, 8 and 9) and pushing the cells to the programmed cell death. Koohpeyma, F et al, demonstrated that treatment with L-carnitine significantly reduced the gene expression of Caspase-9 and significantly increased the gene expression of Bcl-2.²⁵

CONCLUSION

the present work showed that cyclophosphamide treatment produced hazardous effects on the histological structure of the testis. These effects appeared as destruction in the germinal epithelium and sperms with marked reduction in spermatozoa.

These alterations may lead to infertility. Supplementation with Royal jelly or l-carnitine produces relative improvement in the testicular tissue parameters.

Conflict of interest : none

REFERENCES

1. Smart E, Lopes F and Rice S. Chemotherapy drugs cyclophosphamide, cisplatin and doxorubicin induce germ cell loss in an in vitro model of the prepubertal testis. *Sci Rep.* 2018; 8, 1773.
2. Anan HH, Wahba NS, Abdallah M.A, et al. A Histological and immunohistochemical study of cyclophosphamide effect on adult rat testis *Int J Sci Rep.* 2017; 3(2):39-48.
3. Eman ME, Walaa ME, Nadia AE, et al. Histological and immunohistochemical study on the effect of gibberellic acid on the seminiferous tubules of testis of adult albino rat and the possible protective role of grape seeds proanthocyanidin extract. *Tanta Medical Journal.* 2017; 45 (2):79–91.
4. Stukenborg JB, Kjartansdottir KR, Reda A. "Male germ cell development in humans". *Horm Res Paediatr.* 2014; 81(1):2-12.
5. Salazar-Olivo LA and Paz-Gonzalez V. Screening of biological activities present in honeybee (*Apis mellifera*) royal jelly. *Toxicol. In Vitro.* 2005. 19: 645-51.
6. Adewoyin M, Ibrahim M, Roszaman R, et al. Male Infertility: The Effect of Natural Antioxidants and Phytochemicals on Seminal Oxidative Stress. *Diseases.* 2017; 5(1):9.
7. Mongioi L, Calogero AE, Vicari E, et al. The role of carnitine in male infertility. *Andrology.* 2016; 4(5):800–7.
8. Obeys AK, and Mahood AK. Histological study of the effect of Piroxicam on testes of albino mice *Mus musculus.* *J. of university of Anbar for pure science;* 2013; 7(2): 1-11.
9. Stukenborg JB, Kjartansdottir KR, Reda A. "Male germ cell development in humans". *Horm Res Paediatr.* 2014; 81(1):2-12.
10. Fawcett DW, and Jensch RP. Bloom and Fawcett Concise Histology, 2nd ed., Arnold, A member of the Hodder Headline Group, London, New York New Delhi chapter 20. 2002; :270-274.
11. Kamarzaman S, Sha'ban M, and Abdul Rahman S. Effects on Mouse Spermatogenesis and DNA Fragmentation Following Exposure to Cyclophosphamide and Thymoquinone. *European International Journal of Science and Technology.* Vol. 2 No. 2013; 7:1-18.
12. Kanth MA, Kaur P, Ahmad B, et al. Histological Effect of Anticancer Drug Cyclophosphamide (CPA) on Testis of Rattus Rattus. *Indo American*

- Journal of Pharmaceutical Research* 2014; 4 (5):2645-9.
13. Oluwole OF, Salami SA, Ogunwole E, et al. Implication of caffeine consumption and recovery on the reproductive functions of adult male Wistar rats. *J Basic Clin Physiol Pharmacol*, 2016; 27(5): 483-491.
 14. Nie J, and Hou FF. Role of reactive oxygen species in the renal fibrosis. *Chin Med J (Engl)*. 2012; 125 (14): 2598-602.
 15. Karaca T, Demirtaş S, Karaboğa I, et al. Protective effects of royal jelly against testicular damage in streptozotocin-induced diabetic rats. *Turk J Med Sci*. 2015; 45(1):27-32.
 16. Zou W, Yue P, Lin N, He M, and Zhou Z. Vitamin C inactivates the proteasome inhibitor PS-341 in human cancer cells. *Clin.Cancer Res*. 2006; 12: 273-80.
 17. Shi B, Zhao X, Wang C, et al. L-carnitine and fat type in the maternal diet during gestation and lactation modify the fatty acid composition and expression of lipid metabolism-related genes in piglets. *Journal of Animal Physiology and Animal Nutrition (Berl)*. 2019; 103(4), 1207–17.
 18. Zhang X, Cui Y, Dong L, et al. The efficacy of combined l-carnitine and l-acetyl carnitine in men with idiopathic oligoasthenoteratozoospermia: A systematic review and meta-analysis. *Andrologia*. 2020; 52(2): 13470.
 19. Hoda HA, Nashwa SW and Maha AA. Histological and immunohistochemical study of cyclophosphamide effect on adult rat testis. *International Journal of Scientific Reports*. 2017; 3: 39-48.
 20. Fath A, Bethke PC and Jones RL. Enzymes that scavenge reactive oxygen species are down-regulated prior to gibberellic acid-induced programmed cell death in barley aleurone. *Plant Physiol*. 2001; 126(1):156-66.
 21. Roy VK, Verma R, and Krishna A. Carnitine-mediated antioxidant enzyme activity and Bcl2 expression involves peroxisome proliferator-activated receptor- γ coactivator-1 α in mouse testis. *Reproduction, Fertility, and Development*. 2017; 29(6), 1057–63.
 22. Mardanshahi T, Rezaei N, Zare Z, et al. Effects of L-Carnitine on the sperm parameters disorders, apoptosis of spermatogenic cells and testis histopathology in diabetic Rats. *Int J Reprod Biomed*. 2019; 17(5):325.
 23. Vardiyan R, Ezati D, Anvari M, et al. Effect of L-carnitine on the expression of the apoptotic genes Bcl-2 and Bax. *Clin Exp Reprod Med*. 2020; 47(3):155.
 24. Mohammadi V, Sharifi SD, Sharafi M, et al. Dietary L-carnitine affects the expression of genes involved in apoptosis and fatty acid metabolism in rooster testes. *Andrologia*. 2020; 52: 13876.
 25. Koohpeyma F, Siri M and Allahyari S. The effects of L-carnitine on renal function and gene expression of caspase-9 and Bcl-2 in monosodium glutamate-induced rats. *BMC Nephrol*. 2021; 22: 16.